

The 6<sup>th</sup> International Conference on Applied Energy – ICAE2014

## Radiative efficiency estimation of organic substance based on group contribution method

Xinxin Zhang<sup>a,b,c,\*</sup>, Noriyuki Kobayashi<sup>c</sup>, Maogang He<sup>b</sup>, Jingfu Wang<sup>a</sup>

<sup>a</sup>Key Laboratory of Enhanced Heat Transfer and Energy Conservation, Ministry of Education, College of Environmental and Energy Engineering, Beijing University of Technology, Beijing 100124, P R China

<sup>b</sup>Key Laboratory of Thermo-Fluid Science and Engineering, Ministry of Education, School of Energy and Power Engineering, Xi'an Jiaotong University, Xi'an, Shaanxi 710049, P R China

<sup>c</sup>Department of Chemical Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8603, Japan

### Abstract

The development of organic working fluid has entered the environment protection era. Ozone depletion potential (ODP) and global warming potential (GWP) are two most important indices of the organic working fluid. Nowadays, more and more attention has been paid to GWP. The calculation of GWP is an extremely complicated process which involves interactions between surface and atmosphere such as atmospheric radiative transfer and atmospheric chemical reactions. GWP of a substance is related to its atmospheric abundance and is a variable in itself. However, radiative efficiency is an intermediate parameter for GWP calculation and it is a constant value used to describe inherent property of a substance. In this paper, the group contribution method was adopted to estimate the radiative efficiency of the organic substance which contains more than one carbon atom. In most cases, the estimation value and the standard value are in a good agreement and most errors are less than 20%, and the biggest estimation error occurs in the estimation of the radiative efficiency of fluorinated ethers which contains several carbon atoms because of the complicated structure of fluorinated ethers compared with hydrocarbon. This estimation method can be used to predict the radiative efficiency of newly developed organic working fluids and its accuracy can meet the requirement of engineering application.

© 2014 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Peer-review under responsibility of the Organizing Committee of ICAE2014

**Keywords:** Organic substance; Global warming potential; Radiative efficiency; Group contribution method

### 1. Introduction

\* Corresponding author. Tel.: +86-10-6739-1612 ; fax: +86-10-6739-2774 .

E-mail address: [xinxinzhang@bjut.edu.cn](mailto:xinxinzhang@bjut.edu.cn) .

In thermodynamics, there are mainly two applications of organic working fluid. One is used as refrigerant in refrigeration cycle which is the primary application of the organic working fluid. The other is low-grade heat recovery through an organic Rankine cycle(ORC). Indicators for quantitative comparison of the various organic working fluids are ODP(Ozone depletion Potential) and GWP, which are closely related to their atmospheric lifetime(ALT). ODP of one- and two-carbon HFCs(hydrofluorocarbons) and HCFCs((hydrochlorofluorocarbons) can already be estimated with a good precision[1]. However, GWP of organic working fluid cannot be estimated due to its extreme calculation complication. The calculation of GWP involves interactions between surface and atmosphere such as atmospheric radiative transfer and atmospheric chemical reactions. From the definition of GWP given by Intergovernmental Panel on Climate Change(IPCC)[2], it can be seen that GWP of a substance is related to its atmospheric abundance and is a variable in itself. However, radiative efficiency is an intermediate parameter for GWP calculation and it is a constant value used to describe inherent property of a substance. The group contribution method has been widely used and achieved good results in property estimation of organic matter due to its simplicity, reliable accuracy, and wide applicability[3-5]. Therefore, in this paper, the group contribution method was adopted to estimate the radiative efficiency of the organic substance which contains more than one carbon atom. The estimation result offers a good and convenient reference for the development of new organic working fluids.

## 2. Group Division

Molecule was made up of atoms through chemical bond. The existence of intermolecular force makes the distribution of atoms in space not random and homogeneous. The distribution shows a group characteristic via some certain space structures. These groups have an important influence to many properties of substance and this influence remains the same in difference substance. The molecular property is an additive property. The element contribution remains the same in different molecule. In this paper, the group is divided according to the radiative efficiency published by IPCC[2] and the different type of substance. Considering the newly developed organic working fluid usually contains more than one carbon atom, only radiative efficiency of organic matters which contain 2 to 4 carbon atoms are calculated. Table 1 lists the groups and their contribution to the organic matters of different types.

Table 1. Groups and their contribution to the organic matters of different types

Number	Group	Contribution to radiative efficiency of CFC, HCFC, and Halon	Contribution to radiative efficiency of HFC	Contribution to radiative efficiency of fluorinated ethers
1	-CH <sub>3</sub>	0.005	0.015	0.005
2	-CCl <sub>3</sub>	0.015	/	/
3	-CF <sub>3</sub>	0.025	0.115	0.115
4	-CF <sub>2</sub> -	0.06	0.03	0.01
5	-CClF <sub>2</sub>	0.155	/	/
6	-CCl <sub>2</sub> F	0.145	/	/
7	-CBrF <sub>2</sub>	0.165	/	/
8	-CHCl <sub>2</sub>	0.115	/	/
9	-CHClF	0.195	/	/
10	-CH <sub>2</sub> -	/	0.05	0.002
11	-CHF <sub>2</sub>	/	0.095	0.095
12	-CH <sub>2</sub> F	/	0.145	/
13	-CHF-	/	0.03	/
14	-CHCl-	/	/	0.002
15	-O-	/	/	0.23

NOTE: the unit in table is  $W \cdot m^{-2} \cdot ppb^{-1}$

### 3. Result and Discussion

Table 2 lists the estimation results of radiative efficiency of CFC, HCFC, and Halon. Relative error between estimation results and standard value published by IPCC can also be found in Table 2. Table 3 lists the estimation results and relative error of HFC and the corresponding value of fluorinated ethers can be found in Table 4. From these tables, it can be seen that in most cases, the estimation value and the standard value are in a good agreement and most errors are less than 20%, and the biggest estimation error occurs in the estimation of the radiative efficiency of fluorinated ethers which contains several carbon atoms because of the complicated structure of fluorinated ethers compared with hydrocarbon. The estimation accuracy is as fine as that of similar work on estimation of tropospheric lifetimes and ozone-depletion potentials of one- and two-carbon HFC and HCFC by other researchers[1].

Table 2. Estimation results and relative error of CFC, HCFC, and Halon

Working fluid	Chemical formula	Combination of the group( in number)	Standard value of IPCC/ $W \cdot m^{-2} \cdot ppb^{-1}$	Estimation Result/ $W \cdot m^{-2} \cdot ppb^{-1}$	Relative Error/%
CFC-113	$CCl_2FCClF_2$	5+6	0.3	0.3	0
CFC-114	$CClF_2CClF_2$	5+5	0.31	0.31	0
CFC-115	$CClF_2CF_3$	3+5	0.18	0.18	0
Halon-2402	$CBrF_2CBrF_2$	7+7	0.33	0.33	0
Methyl chloroform	$CH_3CCl_3$	1+2	0.06	0.02	66.67
HCFC-123	$CHCl_2CF_3$	3+8	0.14	0.14	0
HCFC-124	$CHClF_2CF_3$	3+9	0.22	0.22	0
HCFC-141b	$CH_3CCl_2F$	1+6	0.14	0.15	7.14
HCFC-142b	$CH_3CClF_2$	1+5	0.2	0.16	20
HCFC-225ca	$CHCl_2CF_2CF_3$	3+4+8	0.2	0.2	0
HCFC-225cb	$CHClF_2CF_2CClF_2$	4+5+9	0.32	0.41	28.13

Table 3. Estimation results and relative error of HFC

Working fluid	Chemical formula	Combination of the group( in number)	Standard value of IPCC/ $W \cdot m^{-2} \cdot ppb^{-1}$	Estimation Result/ $W \cdot m^{-2} \cdot ppb^{-1}$	Relative Error/%
HFC-125	$CHF_2CF_3$	3+11	0.23	0.21	8.70
HFC-134a	$CH_2FCF_3$	3+12	0.16	0.26	62.5
HFC-143a	$CH_3CF_3$	1+3	0.13	0.13	0
HFC-152a	$CH_3CHF_2$	1+11	0.09	0.11	22.22
HFC-227ea	$CF_3CH_2CF_3$	3+3+13	0.26	0.26	0
HFC-236fa	$CF_3CH_2CF_3$	3+3+10	0.28	0.28	0
HFC-245fa	$CHF_2CH_2CF_3$	3+10+11	0.28	0.26	7.14
HFC-365mfc	$CH_3CF_2CH_2CF_3$	1+3+4+10	0.21	0.21	0

Table 4. Estimation results and relative error of fluorinated ethers

Working fluid	Chemical formula	Combination of the group( in number)	Standard value of IPCC/ $W \cdot m^{-2} \cdot ppb^{-1}$	Estimation Result/ $W \cdot m^{-2} \cdot ppb^{-1}$	Relative Error/%
HFE-125	$CHF_2OCF_3$	3+11+15	0.44	0.44	0
HFE-134	$CHF_2OCHF_2$	11+11+15	0.45	0.42	6.67
HFE-143a	$CH_3OCF_3$	1+3+15	0.27	0.35	29.63
HCFE-235da2	$CHF_2OCHClCF_3$	3+11+14+15	0.38	0.442	16.32
HFE-245cb2	$CH_3OCF_2CHF_2$	1+4+11+15	0.32	0.34	6.25
HFE-245fa2	$CHF_2OCH_2CF_3$	3+10+11+15	0.31	0.442	42.58
HFE-254cb2	$CH_3OCF_2CHF_2$	1+4+11+15	0.28	0.34	21.43
HFE-347mcc3	$CH_3OCF_2CF_2CF_3$	1+3+4+4+15	0.34	0.37	8.82
HFE-347pcf2	$CHF_2CF_2OCH_2CF_3$	3+4+10+11+15	0.25	0.452	80.8
HFE-356pcc3	$CH_3OCF_2CF_2CHF_2$	1+4+4+11+15	0.93	0.35	62.37

#### 4. Conclusion

The group contribution method was adopted to estimate the radiative efficiency of the organic substance which contains more than one carbon atom. In most cases, the estimation value and the standard value are in a good agreement and most errors are less than 20%, and the biggest estimation error occurs in the estimation of the radiative efficiency of fluorinated ethers which contains several carbon atoms because of the complicated structure of fluorinated ethers compared with hydrocarbon. This estimation method can be used to predict the radiative efficiency of newly developed organic working fluids and its accuracy can meet the requirement of engineering application.

#### Acknowledgements

This work was done in China and Japan with the support from the National Key Basic Research Program of China (973 Program, No.2013CB228306) and China Scholarship Council (CSC) respectively. The authors gratefully acknowledge them for financial support of this work.

#### References

- [1] Nlmltz JS, Skaggs SR. Estimating tropospheric lifetimes and ozone-depletion potentials of one- and two-carbon hydrofluorocarbons and hydrochlorofluorocarbons. *Environ Sci Technol* 1992;**26**:739-744.
- [2] Intergovernmental Panel on Climate Change. *Climate Change 2007 The Physical Science Basis, Working Group I Contribution to the Fourth Assessment Report of the IPCC*. Cambridge: Cambridge University Press; 2007.
- [3] Khetib Y, Meniai AH, Lallemand A. Computer-aided design of CFC and HCFC substitutes using group contribution methods. *Desalination* 2009;**239**:82-91.
- [4] Ismail HM, Samir BB, El-Harbawi M. Development of a novel mathematical model using a group contribution method for prediction of ionic liquid toxicities. *Chemosphere* 2011;**85**:990-4.
- [5] Ceriani R, Gani R, Meirelles AJA. Prediction of heat capacities and heats of vaporization of organic liquids by group contribution methods. *Fluid Phase Equilibria* 2009;**28**:49-55.



#### Biography

Dr. Xinxin Zhang graduated from Xi'an Jiaotong University and was a joint training doctoral student of Nagoya University supported by China Scholarship Council. Now he works in Beijing University of Technology and his research interest is new-type and combined thermodynamic cycle and waste heat recovery through thermodynamic approach.